

Histopathological study of *Capparis spinosa* on the healing of experimental Achilles tendon injury in rabbits

Hamid Reza Moslemi^{1*}; Seyyed Morteza Haeryan²; Meysam Tehrani Sharif³
and Ramin Mazaheri Nezhad Fard⁴

¹ Assistant Professor, Department of Clinical Sciences, Faculty of Veterinary Medicine, University of Semnan, Semnan, Iran

² DVM Graduated, Faculty of Veterinary Medicine, Garmsar Branch, Islamic Azad University, Garmsar, Iran

³ Assistant Professor, Department of Clinical Sciences, Faculty of Veterinary Medicine, Garmsar Branch, Islamic Azad University, Garmsar, Iran

⁴ Assistant Professor, Department of Pathobiology, Faculty of Public Health and Food Microbiology Research Center, Tehran University of Medical Sciences, Tehran, Iran

Received: 10.12.2019

Accepted: 11.06.2020

Abstract

The critical role of tendons in body mechanics and injury and degeneration of this tissue can highly be debilitating, resulting in substantial pain, disability and costs. *Capparis spinosa* is one of the most common aromatic plants growing in Iran. The major objective of the study was to assess Achilles tendon healing in a rabbit model by local injection of ethanolic extract of *C. spinosa*. Nine adult white New Zealand male rabbits were anesthetized and partial thickness tenotomies were created on both hindlimbs. The *C. spinosa* extract and normal saline were respectively injected daily to the treatment and control groups for three days post-operatively. Histological analysis on days 7, 14 and 28 post-rupture demonstrated higher regenerating activity and capacity in treated groups than the control group. This was illustrated by fewer inflammatory cells, a larger number of blood vessels, further fibroblasts and increased structural organization with further longitudinally oriented collagen fibers in the treated group. In summary, these results suggest that use of *C. spinosa* extract can promote the healing process of damaged Achilles tendons in rabbits.

Key words: *Capparis spinosa*, Achilles tendon, Rabbit

Introduction

Tendon is a compositionally complex tissue with a predominantly mechanical function of translating muscular contractions into joint movement by transmitting forces from muscles to bones. Tendons are able to store elastic energy and withstand high tensile forces; on which, body locomotion is entirely dependent (Aslan et al. 2008). Damages to these structures affect the natural balance between stability and mobility; thus, altering joint kinematics and ultimately lead

to destruction of the joints (Ferrara 2002). Owing to critical roles of this tissue in body mechanics, injury and degeneration of tendons can highly be debilitating, which can result in substantial pain, disability and health-care costs. Delayed tendon healing and adhesion are still complications that occur most often after tendon repairs. Following tendon injuries, process of healing or tissue repair starts that can largely be divided into three overlapping phases of inflammatory, repairing and

* **Corresponding Author:** Hamid Reza Moslemi, Assistant Professor, Department of Clinical Sciences, Faculty of Veterinary Medicine, University of Semnan, Semnan, Iran, E-mail: h.moslemi@semnan.ac.ir



© 2020 by the authors. Licensee SCU, Ahvaz, Iran. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0 license) (<http://creativecommons.org/licenses/by-nc/4.0/>).

remodeling (Bosch et al. 2011, Kuo et al. 2003, Tohidnezhad et al. 2011). Antioxidants play key roles in healing process of each phase (Martin 1996). The process of inflammation normally leads to release of biologically active mediators to attract neutrophils, leukocytes and monocytes to the wound area to attack foreign debris and microorganisms through phagocytosis. This process results in production of oxygen-free radicals such as hydrogen peroxide, superoxide anion and hydroxyl anion. Excesses in these free radicals cause tissue damages in humans or animals if they overwhelm natural antioxidant enzymes of the host such as catalase, superoxide dismutase and glutathione peroxidase. Therefore, antioxidants prevent activity of the free radicals and thereby prevent damages to cells and tissues, provide protection to humans and animals and enhance healing of infected and non-infected wounds (Houghton et al. 2005, Martin 1996).

Capparis spinosa or caper (Family Capparidaceae) is one of the most common aromatic plants growing wild in dry regions around the West and Central Asia and the Mediterranean basin. Caper has been known for centuries in traditional phytomedicine, which has described its properties for several purposes. Studies have revealed that *Capparis* includes anti-oxidative (Mousavi et al. 2016), anti-fungal (Shtayeh and Abu Ghdeib 1999), anti-inflammatory (Al-Said et al. 1999), anti-diabetic (Ziyyat et al. 1997) and anti-hyperlipidemic (Eddouks et al. 2005) activities and contains constituents of polyherbal formulations to treat liver ailments (Madrigal-Santillan et al. 2014), diuretic, antihypertensive and poultice (Calis et al. 1999). Various parts of the caper plant that can be used as drugs, cosmetics and foods are further used in various regions for landscaping, control of erosion and animal feeding (Ozcan and Akgul 1998). Based on the extensive anti-inflammatory effects of *C. spinosa* and

effects of inflammatory control on tendon repair processes, the use of this plant is suggested to play effective roles in tendon repair processes. Therefore, the current study investigated the healing effects of local injections of *C. spinosa* extract on experimental Achilles tendon injuries in rabbits.

Materials and Methods

Animals

This study was carried out on nine adult white New Zealand male rabbits with a mean bodyweight of 2–2.5 Kg. Before beginning of the experiment, rabbits were housed for two weeks to adapt to the environment. The animals were maintained under controlled conditions of 25 °C ±1 and 12 h light-dark cycles and had free access to standard chow diet and water throughout the study. The study was approved by the Institutional Animal Care and Use and Committee and carried out based on the guidelines from the National Institute of Health (NIH).

Plant materials and extract preparation

Roots of *C. spinosa* were collected from Varamin, Province of Tehran, Iran. Specimens were deposited at the herbarium of the institute of medicinal plants, Karaj, Iran. Roots of the plant were dried and grounded into fine powder using electric blender. The extract was prepared using cold maceration with distilled water for 24 h. The plant powder (50 gr) was suspended at 100 ml of ethanol for 24 h at room temperature. The mixture was filtered using fine muslin cloths followed by Whatman no. 1 filter papers. Extract was concentrated using vacuum distillation (Ezatpour et al. 2015).

Experimental protocol

Rabbits were anesthetized with ketamine hydrochloride (5%, 35 mg/kg) in combination with xylazine (2%, 5 mg/kg) intramuscularly (Fazel and Moslemi 2017). The anesthesia was maintained with

inhalation isoflurane (2.5%). Surgery was carried out on both hindlimbs; with left hindlimb served as control (Fazel and Moslemi 2017). The surgical area was prepared aseptically. A longitudinal skin incision was made over the Achilles tendon and the paratenon was exposed and incised longitudinally as a separate layer. The three bundles of Achilles tendon were identified and the central bundle was separated bluntly. A partial-thickness tenotomy (approximately 50% of the tendon bundle width with 1 cm length) was created at 2 cm to the proximal side of the Achilles tendon-calcaneus junction. The severed tendon was not sutured. The partial tenotomy allowed the rest of the tendon to act as an internal splint for the non-immobilized repair (Fazel and Moslemi 2017). One milliliter of the *C. spinosa* extract and normal saline were injected daily to the treatment and control groups respectively for three days post-operatively. After the surgery, rabbits were recovered from the anesthesia in a warm recovery chamber under continuous observation. Following recovery, animals

were returned to the individual cages for the rest of the experiment. Flunixin (Razak Co. Iran) as analgesic (2.5 mg/kg IM) and enrofloxacin as antibiotic (5 mg/kg, IM) agents were administered to rabbits one hour pre-operation and were continued for three days.

Histopathological studies

Animals were postoperatively euthanized on Days 7, 14 and 28 using an overdose of thiopental sodium. Then, Achilles tendons were removed by dissection. Number of specimens was three per day. Specimens were fixed in 10% buffered formalin, dehydrated and embedded in paraffin. After 10 days, longitudinal sections of the tendon were stained with hematoxylin and eosin (H&E). Stained specimens were microscopically studied to assess extent and severity of inflammation, angiogenesis, fibroplasia and complete tendon healing, which were scaled from 0 to 3 by defined criteria (Hadjipour et al. 2008). These criteria are shown in Table 1.

Table 1. Grading system for the histopathological study

Tendon repair assessment score				
	0	1	2	3
Extent and severity of the inflammation	Inflammatory cells were not seen	Observation of inflammatory cells at two microscopic fields	Observation of inflammatory cells at 3–5 microscopic fields	Observation of inflammatory cells at more than five microscopic fields
Angiogenesis	Blood vessels were not seen	Existing of 0–2 blood vessels	Existing of 3–4 blood vessels	Existing of more than four blood vessels
Fibroplasia	Recording of few thin collagen fibers with numerous fibroblasts	Recording of thin collagen fibers with very numerous fibroblasts	Recording of thick collagen fibers with numerous fibroblasts	Recording of abundant thick collagen fibers with few fibroblasts
Complete tendon healing	Observation of inflammatory cells; no observation of blood vessels, fibroblasts or collagen fibers	Contemporary observation of inflammatory cells, blood vessels, fibroblasts and collagen fibers	Contemporary observation of blood vessels, fibroblasts and collagen fibers; no observation of inflammatory cells	Observation of fibroblasts, thick and compact collagen bundle; no observation of inflammatory cells or blood vessels

Data analysis

Results were expressed as mean±SD (standard deviation). Significant differences between groups were assessed using Mann-Whitney U-test and SPSS Software v.16.0 (SPSS, USA). Differences were considered significant when $P < 0.05$.

Results

The average scores of histopathological changes of the complete tendon healing in treatment and control groups are shown in Table 2. In the present study, number of fibroblasts and degree of neovascularization were higher in *C. spinosa* treated group than in control group on Days 7 and 14 of injury. In fibroplasia and angiogenesis on Day 7, mean±SD of *C. spinosa* treated group and control group were 1.66 ± 0.57 and 1, respectively. These factors on day 14 were 2.33 ± 0.57 and 1.66 ± 0.57 , respectively. Based on these findings, differences on day 7 were significant ($P < 0.05$) while differences on day 14 were not significant statistically ($P \geq 0.05$). This result has suggested that *C. spinosa* promotes fibroblast migration and/or proliferation as well as neovascularization at early stages of tendon healing. At these times, injured tendons in treatment and control groups showed structural organizations of the new collagen fibers aligning with the functional loading axis in a further pronounced manner in *C. spinosa* treated group. This exhibited hypocellular areas of further disorganized collagens; for example, collagen fibers diverging from the functional axis (Fig. 1 and 2A, B). On day 7, the average score of histopathological changes of the complete tendon healing in *C. spinosa* treated group and the control group were respectively 1.66 ± 0.57 and 1 with significant differences ($P < 0.05$). On day 14, the

average score of histopathological changes of the complete tendon healing in *C. spinosa* treated group and control group were respectively 2.33 ± 0.57 and 1.66 ± 0.57 with no statistically significant differences ($P > 0.05$). On day 28 for fibroplasia and angiogenesis, the mean±SD of *C. spinosa* treated group and control group were respectively 2.66 ± 0.57 and 2 with statistically significant references ($P < 0.05$). The number of fibroblast decreased in *C. spinosa* treated group at day 28. In consistency with these changes, thick collagen fiber bundles were aligned in one direction, parallel to the long axis of the tendon. However, the control group showed deposition of thin collagen fibers and hypercellularity characterized by increased fibroblasts (Fig. 3A, B). On day 28, the average score of histopathological changes of the complete tendon healing in *C. spinosa* treated group and control group were respectively 2.66 ± 0.57 and 2 with statistically significant differences ($P < 0.05$). Furthermore, results of various days from each group were compared to each other with differences not significant statistically ($P > 0.05$).

Table 2. The average score of histopathological changes of the complete tendon healing in the two groups

Day	<i>C. spinosa</i> extract	Control
7	1.66 ± 0.57 ^{A,a}	1 ^{B, b}
14	2.33 ± 0.57 ^A	1.66 ± 0.57 ^B
28	2.66 ± 0.57 ^{A,a}	2 ^{B, b}

No significant differences were seen between similar capital letters in each column while significant differences were observed between the non-identical lowercases in each row. Data were represented as mean±SD with significant differences ($P < 0.05$).




Fig. 1. Photomicrograph of the injured tendon in control group (A) and *C. spinossa* treated group (B) on day 7 post-operation. Note less hypercellularity, fewer new blood vessels and better healing process in treated group (H&E, $\times 200$)




Fig. 2. Photomicrograph of the injured tendon in control group (A) and *C. spinossa* treated group (B) on day 14 post-operation. Note less hypercellularity and more mature fibrous connective tissue and better healing process in *C. spinossa* treated group (H&E, $\times 200$)




Fig. 3. Photomicrograph of the injured tendon in control group (A) and *C. spinossa* treated group (B) on day 28 post-operation. Note thick collagen fibers in parallel arrangement in treatment group. However, the control group showed deposition of thin collagen fibers with blood vessels and hypercellularity (H&E, $\times 200$)

Discussion

Injuries and degenerative conditions of tendons represent almost 50% of the musculoskeletal injuries treated in

orthopedic clinics (Schweitzer et al. 2010). Similar to other connective tissues, the tendon repair process has been an attractive

subject for the researchers for many years. It is well known that increased blood supply enhances repair process in all types of connective tissues (Fazel and Moslemi 2017, Sharma and Maffulli 2005). The major objective of the current study was to improve and accelerate Achilles tendon healing in a rabbit model using local injection of the ethanolic extract of *C. spinosa*. The histological analysis on days 7, 14 and 28 post-rupture demonstrated a higher regenerating activity and capacity in treated groups than control group. This was illustrated by fewer inflammatory cells, larger numbers of blood vessels, more fibroblasts and increased structural organizations with more longitudinally oriented collagen fibers in treated groups. These results suggested that the *C. spinosa* treatment promoted tendon tissue repair.

Healing procedures usually include suppression of inflammation, cell proliferation and contraction of the collagen tissues; therefore, these processes can be delayed by reactive oxygen species or microbial infections (Houghton et al. 2005, Velnar et al. 2009). Tendon injuries have been shown to benefit from antioxidant therapy (Park et al. 2010). Antioxidant and free-radical scavenging activities (Kordali et al. 2005, Lopes-Lutz et al. 2008, Taherkhani et al. 2013) and anti-inflammatory activity (Kordali et al. 2005) have been reported for the essential oils of *C. spinosa*. Plant-derived antioxidants such as phenolic acids, flavonols and flavones can postpone or prevent invasion of degenerative diseases because of their redox properties, which allow them to be active as reducing agents, hydrogen donors, hydroxyl radicals (OH) or superoxide radical (O₂) scavengers (Goorani et al. 2019, Govindarajan et al. 2005). Antioxidants enhance healing of infected and non-infected wounds by reducing damages caused by oxygen radicals (Houghton et al. 2005).

Total phenolic content of the extract often determines its pharmacological

effects such as antioxidant activity. Components other than phenolic or flavonoid contents may play important roles in its antioxidant activity. Studies have reported that phenolic compounds display antioxidant activity as a result of their capacity to scavenge free-radicals. Phenolic compounds can act as antioxidants by chelating metal ions, preventing radical formation and improving antioxidant endogenous system (Seyoum et al. 2006, Qingming et al. 2010). Earlier studies have shown that flavonoids include direct roles in scavenging ROS, which can reverse lipid oxidation *in vitro* and improve body antioxidant enzyme activity and decrease peroxide formation *in vivo* (Nakao et al. 2011). Rutin, quercetin, 3-rutinoside, keampferol 3-rutinoside and kaempferol 3-rhamnosylrutinoside as phenolic compounds have been reported in *C. spinosa* (Tlili et al. 2011).

Various mechanisms, including reducing capacity, prevention of chain initiation, binding of transition metal ion catalysts, decomposition of peroxides, prevention of continued hydrogen abstraction and radical scavenging, have been suggested to explain antioxidant activities (Braca et al 2003, Re et al. 1999). The reducing capacity of a compound may serve as a significant indicator of its potential antioxidant activity. Extracts of *C. spinosa* exhibit effective reducing capacities at all concentration points. Reducing capacity of the extracts increased with increased concentration. The reducing properties are generally associated with the presence of reductones (Duh 1998), which have been shown to include antioxidant activities by breaking the free radical chain and donating hydrogen atoms. Reductones are reported to react with certain precursors of peroxide; thus, preventing peroxide formation (Li et al. 2007). Therefore, it seems that the phenolic constituents in *C. spinosa* are responsible for its antioxidant and free radical-scavenging activities. Significant differences between the treatment and

control groups in the current study suggest that *C. spinosa* is a good source of phenolic compounds, especially flavonoids. Despite high antioxidant activity of this plant and its effects on tendon healing, antioxidant

assessment was not carried out in this study. Results of the study have demonstrated that *C. spinosa* includes potentials to improve the healing process of damaged Achilles tendons.

Acknowledgement

The authors would like to thanks Mr. K. Nematollahi for technical support.

Conflict of interest

The authors have no conflicts of interest to declare.

Funding

The authors received no funding for this work.

References

- Al-Said, M.S.; Abdelsattar, E.A.; Khalifa, S.I. and El-Ferly, F.S. (1999). Isolation and identification of an anti-inflammatory principle from *Capparis spinosa*. *Die Pharmazie*. 43: 640-641.
- Aslan, H.; Kimelman-Bleich, N.; Pelled, G. and Gazit, D. (2008). Molecular targets for tendon neof ormation. *Journal of Clinical Investigation*, 118: 439-444.
- Bosch, G.; Moleman, M.; Barneveld, A.; Van Weeren, P.R. and Van Schie, H.T. (2011). The effect of platelet-rich plasma on the neovascularization of surgically created equine superficial digital flexor tendon lesions. *Scandinavian Journal of Medicine Science In Sports*. 21: 554-561.
- Braca, A.; Fico, G.; Morelli, I.; Simone, F.; Tom, F. and De Tommasi, N. (2003). Antioxidant and free radical scavenging activity of flavonoid glycosides from different *Aconitum* species. *Journal of Ethnopharmacology*. 86(1): 63-67.
- Çalis, I.; Kuruüzüm, A. and Rüedi, P. (1999). 1H-indole-3-acetonitrile glycosides from *Capparis spinosa* fruits. *Phytochemistry*. 50: 1205-1208.
- Duh, P.D. (1998). Antioxidant activity of burdock (*Arctium lappa* Linne): its scavenging effect on free radical and active oxygen. *Journal of the American Oil Chemists' Society*. 75: 455-461.
- Eddouks, M.; Lemhadri, A. and Michel, J.B. (2005). Hypolipidemic activity of aqueous extract of *Capparis spinosa* L. in normal and diabetic rats. *Journal of Ethnopharmacology*. 98(3): 345-50.
- Ezatpour, B., Saedi Dezaki, E., Mahmoudvand, H., Azadpour, M. and Ezzatkah, F. (2015). *In vitro* and *in vivo* antileishmanial effects of *Pistacia khinjuk* against *Leishmania tropica* and *Leishmania major*. Evidence Based Complementary and Alternative Medicine, Article ID 149707, <http://dx.doi.org/10.1155/2015/149707>
- Fazel, H. and Moslemi, H.R. (2017). Antioxidant properties of *Pistacia khinjuk* accelerate healing of the experimental achilles tendon injury in rabbits. *Journal of Medicinal Plants*. 16(63): 33-42.
- Ferrara, N. (2002). Role of vascular endothelial growth factor in physiologic and pathologic angiogenesis: therapeutic implications. *Seminars In Oncology*, 29: 10-14.
- Goorani, S.; Zangeneh, M.; Koohi, M. and Seydi, N. (2019). Assessment of antioxidant and cutaneous wound healing effects of *Falcaria vulgaris* aqueous extract in Wistar male rats. *Comparative Clinical Pathology*, 28(2): 435-445.
- Govindarajan, R.; Vijayakumar, M. and Pushpangadan, P. (2005). Antioxidant approach to disease management and the role of 'Rasayana' herbs of Ayurveda. *Journal of Ethnopharmacology*, 99: 165-178.
- Hadjipour, N.; Marjani, M.; Sedaghat, R.; Marjanmehr, SH. (2008). Histopathological Absorption Trend Comparison between Chromic Catgut Suture Materials Produced by Different Companies in Rabbits. *Journal of Animal and Veterinary Advances*. 11: 1341-1344.
- Houghton, P.J.; Hylands, P.J.; Mensah, A.Y.; Hensel, A. and Deters, A.M. (2005). *In vitro* tests and ethnopharmacological investigations: wound healing as an example. *Journal of Ethnopharmacology*, 100(1-2): 100-107.

- Kordali, S.; Cakir, A.; Mavi, A.; Kilic, H. and Yildirim, A. (2005). Screening of chemical composition and antifungal and antioxidant activities of the essential oils from three Turkish *Artemisia* species. *Journal of Agricultural and Food Chemistry*, 53(5): 1408-1416.
- Kuo, Y.R.; Kuo, M.H.; Chou, W.C.; Liu, Y.T.; Lutz, B.S. and Jenf, S.F. (2003). One-stage reconstruction of soft tissue and Achilles tendon defects using a composite free anterolateral thigh flap with vascularized fascia lata: clinical experience and functional assessment. *Annals of Plastic Surgery*, 50: 149.
- Li, X.M.; Li, X.L. and Zhou, A.G. (2007). Evaluation of antioxidant activity of the polysaccharides extracted from *Lycium barbarum* fruits in vitro. *European Polymer Journal*, 43: 488-497.
- Lopes-Lutz, D.; Alviano, D.S.; Alviano, C.S. and Kolodziejczyk, P.P. (2008). Screening of chemical composition, antimicrobial and antioxidant activities of *Artemisia* essential oils. *Phytochemistry*, 69(8): 1732-1738.
- Madrigal-Santillan, E.; Madrigal-Bujaidar, E.; Alvarez-Gonzalez, I.; Sumaya-Martínez, M.T.; Gutierrez-Salinas, J.; Bautista, M. and et al. (2014). Review of natural products with hepatoprotective effects. *World Journal of Gastroenterology*, 20(40): 14787-14804.
- Martin, A. (1996). The use of antioxidants in healing. *Dermatologic Surgery*, 22(2): 156-160.
- Mousavi, S.; Hosseini, A.; Bakhtiari, E. and Rakhshandeh, H. (2016). *Capparis spinosa* reduces Doxorubicin-induced cardio-toxicity in cardiomyoblast cells. *Avicenna Journal of Phytomedicine*, 6: 488-494.
- Nakao, K.; Murata, K.; Itoh, K.; Hanamoto, Y.; Masuda, M.; Moriyama, K. et al. (2011). Anti-hyperuricemia effects of extracts of immature *Citrus unshiu* fruit. *Journal of Traditional Medicines*, 28(1): 10-15.
- Ozcan, M. and Akgul, A. (1998). Influence of species, harvest date and size on composition of capers (*Capparis* spp.) flower buds. *Molecular Nutrition and Food Research*, 42(2): 102-105.
- Park, H.B.; Hah, Y.S.; Yang, J.W.; Nam, J.B.; Cho, S.H. and Jeong, S.T. (2010). Antiapoptotic effects of anthocyanins on rotator cuff tenofibroblasts. *Journal of Orthopaedic Research*, 28: 1162-1169.
- Qingming, Y.; Xianhui, P.; Weibao, K.; Hong, Y.; Yidan, S.; Li, Z. et al. (2010). Antioxidant activities of malt extract from barley (*Hordeum vulgare* L.) toward various oxidative stress *in vitro* and *in vivo*. *Food Chemistry*, 118: 84-89.
- Re, R.; Pellegrini, N.; Protoggente, A.; Pannala, A.; Yang, M. and Rice-Evans, C. (1999). Antioxidant activity applying an improved ABTS radical cation decolorisation assay. *Free Radical Biology and Medicine*, 26: 1231-1237.
- Schweitzer, R.; Zelzer, E. and Volk, T. (2010). Connecting muscles to tendons: tendons and musculoskeletal development in flies and vertebrates. *Development*, 137: 2807-2817.
- Seyoum, A.; Asres, K. and El-Fiky, F.K. (2006). Structure-radical scavenging activity relationships of flavonoids. *Phytochemistry*, 67: 2058-2070.
- Sharma, P. and Maffulli, N. (2005). Tendon injury and tendinopathy: healing and repair. *Journal of Bone and Joint Surgery, American Volume*, 87: 187-202.
- Shtayeh, A. and Abu Ghdeib, S.L. (1999). Antifungal activity of plant extracts against dermatophytes. *Mycoses*, 42: 665-672.
- Taherkhani, M.; Rustaiyan, A. and Taherkhani, T. (2013). Chemical composition, antimicrobial activity, antioxidant and total phenolic content within the leaves essential oil of *Artemisia absinthium* L. growing wild in Iran. *African Journal of Pharmacy and Pharmacology*, 7(2): 30-36.
- Tlili, N.; Nasri, N.; Khaldi, A.; Triki, S. and Munne-Bosch, S. (2011). Phenolic compounds, tocopherols, carotenoids and vitamin C of commercial caper. *Journal of Food Biochemistry*, 35(2): 472-483.
- Tohidnezhad, M.; Varoga, D.; Wruck, C.J.; Brandenburg, L.O.; Seekamp, A.; Shakibaei, M. et al. (2011). Platelet-released growth factors can accelerate tenocyte proliferation and activate the anti-oxidant response element. *Histochemistry and Cell Biology*, 135: 453-460.
- Velnar, T.; Bailey, T. and Smrkolj, V. (2009). The wound healing process: an overview of the cellular and molecular mechanisms. *Journal of International Medical Research*, 37(5): 1528-1542.
- Ziyyat, A.; Legssyer, A.; Mekhfi, H.; Dassouli, A.; Serhouchni, M. and Benjelloun, W. (1997). Phytotherapy of hypertension and diabetes in oriental Morocco. *Journal of Ethnopharmacology*, 58: 45-54.